

Another homogeneous, non-bihomogeneous Peano continuum

Greg Kuperberg

1997

Abstract

K. Kuperberg found a locally connected, finite-dimensional continuum which is homogeneous but not bihomogeneous. We give a similar but simpler example.

In 1988, K. Kuperberg [4] found a locally connected, finite-dimensional continuum which is homogeneous but not bihomogeneous. The basic idea of this construction is to use the fact that a local homeomorphism of $\mu^1 \times \mu^1$, where μ^1 is the Menger curve, either preserves horizontal and vertical fibers or switches them. This places strong restrictions on the possible homeomorphisms of more complicated continua that resemble $\mu^1 \times \mu^1$. Since μ^1 is homogeneous, the restrictions do not necessarily preclude homogeneity.

In this note, we give a similar but slightly simpler example of a locally connected continuum (a Peano continuum) which is homogeneous but not bihomogeneous. The author found the example after some informative discussions with K. Kuperberg.

1 Terminology

A space is *homogeneous* if for every x and y , there is a homeomorphism sending x to y . It is *bihomogeneous* if there is a homeomorphism exchanging x and y . It is *strongly locally homogeneous* if for every x and every open $U \ni x$, there is an open V such that $x \in V \subset U$ and for every $y \in V$, there is a homeomorphism taking x to y which is the identity outside U . Finally, a space is *weakly 2-homogeneous* if for every pair of unordered pairs of distinct points $\{x_1, x_2\}$ and $\{y_1, y_2\}$, there is a homeomorphism taking $\{x_1, x_2\}$ to $\{y_1, y_2\}$.

2 The example

Given a closed PL manifold M of dimension at least $2n + 1$, let $X_0 = M$, and for each $i > 0$, let X_i be a regular neighborhood in X_{i-1} of the n -skeleton of a triangulation of X_{i-1} . The intersection $\mu_M^n = \bigcap_i X_i$ is a locally $n-1$ -connected continuum called a Menger manifold whose homeomorphism type does not depend on the particular triangulations chosen [2]. The homeomorphism type of μ_M^1 also does not depend on M and is therefore that of μ^1 , the Menger curve [1]. The space μ_M^n is strongly locally homogeneous. In general $\pi_k(\mu_M^n) \cong \pi_k(M)$ for $k < n$. The manifold $\mu_{S^{2n+1}}^n$ is simply written as μ^n ; any $x \in \mu_M^n$ has a closed neighborhood homeomorphic to μ^n .

Consider the 21-element $G = \mathbb{Z}/3 \rtimes \mathbb{Z}/7$ which is a non-trivial semidirect product. The group G is generated by elements a and b with relations $a^3 = b^7 = 1$ and $aba^{-1} = b^2$. It has no automorphism sending a to a^{-1} . Let W be a closed, PL 5-manifold with fundamental group G . (One can make a closed, PL 5-manifold whose fundamental group is an arbitrary finitely presented group by first making a 2-complex from the presentation, embedding it in $\mathbb{R}^5$, and gluing two copies of a regular

neighborhood along the boundary.) Let $\widetilde{W}$ be the connected covering of W corresponding to the subgroup generated by b , which is a regular 3-fold covering. Then $\pi_1(\widetilde{W}) \cong \mathbb{Z}/7$, $\mu_{\widetilde{W}}^2$ and μ_W^2 have the same fundamental groups as $\widetilde{W}$ and W , and $\mu_{\widetilde{W}}^2$ is a regular 3-fold covering of μ_W^2 . Evidently, $\mu_{\widetilde{W}}^2$ has a homeomorphism ϕ (a deck translation) of order three with no fixed points which squares elements of its fundamental group.

The example X is a twisted bundle with fiber $\mu_{\widetilde{W}}^2$ and base μ^1 . Specifically, let $f : \mu^1 \rightarrow C$ be a retraction onto a circle C contained in μ^1 , let X' be a $\mu_{\widetilde{W}}^2$ bundle over C with monodromy ϕ , and let X be the pullback of X' under f . Alternatively, μ^1 has a regular covering of itself of order 3 with deck translation α ; let X be the quotient of $\mu_{\widetilde{W}}^2 \times \mu^1$ by the group action $(x, y) \mapsto (\phi(x), \alpha(y))$.

Theorem 1 *The continuum X is homogeneous but not bihomogeneous.*

PROOF: To establish homogeneity, it suffices to note that X is a regular 3-fold covering of $\mu_W^2 \times \mu^1$. Let π be the covering map. Since μ_W^2 and μ^1 are strongly locally homogeneous and compact, for every $x, y \in \mu_W^2 \times \mu^1$, there is a homeomorphism of $\mu_W^2 \times \mu^1$ which takes x to y and which lifts to X . Therefore any fiber of π can be taken to any other by a homeomorphism, and it remains to show that if x and y are in the same fiber, there is a homeomorphism taking x to y . Indeed, there is a deck translation with this property.

To show that X is not bihomogeneous, we define a horizontal fiber of X to be the preimage of a point of $f \circ \pi$ and a vertical fiber to be the preimage of $g \circ \pi$, where f and g are the projections of $\mu_W^2 \times \mu^1$ onto the first and second Cartesian factors, respectively. By a lemma of K. Kuperberg [3, lemma 1], arbitrarily small subsets of $\mu^2 \times \mu^1$ have the property that every open embedding into must take horizontal fibers (meaning $\mu^2 \times \{p\}$) into horizontal fibers and vertical fibers (meaning $\{q\} \times \mu^1$) into vertical fibers. Therefore a homeomorphism ψ of X must preserve horizontal and vertical fibers. In particular, ψ must preserve π -fibers, because they are characterized as the intersections of horizontal and vertical fibers.

We claim that if x and y lie in the same π -fiber, ψ cannot exchange them. Otherwise, ψ fixes the third point of the same fiber. The map ψ restricts to the vertical fiber containing x and y . Since it preserves π -fibers, it further descends to a homeomorphism of a copy of μ_W^2 . Call this homeomorphism ψ also. Since ψ reverses the cyclic ordering of a π -fiber over this copy of μ_W^2 , the induced map on $\pi_1(\mu_W^2) \cong G$ sends a to a^{-1} . But G has no such automorphism. $\square$

3 Remarks and open problems

Clearly, the 21-element group in the above construction can be replaced by many other finite groups. For any $k \geq 1$, there is also a locally k -connected example which is a μ_W^r bundle over $\mu_{S^{2s} \times S^1}^s$ for any $r, s > k$. But if $r = s = 1$, the corresponding bundle becomes bihomogeneous.

Say that a topological space is *primitively homogeneous* if no non-trivial equivalence relation on its points is invariant under all homeomorphisms. (An equivalence relation is *invariant* if that $x \sim y \iff f(x) \sim f(y)$. This is analogous to the notion of a primitive permutation group.) The continuum $\mu^1 \times \mu^1$ is primitively homogeneous and bihomogeneous, but not weakly 2-homogeneous. On the other hand, if one takes a path-connected component of a solenoid (a solenoid composant), removes a point, and takes a path-connected component of the remainder, the result is a topological space embeddable in $\mathbb{R}^n$ which is primitively homogeneous and even weakly 2-homogeneous, but not bihomogeneous.

Question 1 *Is there a continuum which is primitively homogeneous but not bihomogeneous? Primitively homogeneous but neither bihomogeneous nor weakly 2-homogeneous?*

References

- [1] R. D. Anderson. A characterization of the universal curve and the proof of its homogeneity. *Ann. of Math.*, 67:313–324, 1958.
- [2] M. Bestvina. Characterizing k -dimensional universal Menger compacta. *Mem. Amer. Math. Soc.*, 71, 1988.
- [3] K. Kuperberg. Bihomogeneity and Menger manifolds. Preprint.
- [4] K. Kuperberg. On the bihomogeneity problem of Knaster. *Trans. Amer. Math. Soc.*, 321:129–143, 1990.